

VTE Prophylaxis Options for Surgery	
Surgery Type	Recommended Prophylaxis Options¹
Intracranial Neurosurgery Appendix A, Table 5.17	Any of the following: <ul style="list-style-type: none"> • Intermittent pneumatic compression devices (IPC) with or without graduated compression stockings (GCS) • Low-dose unfractionated heparin (LDUH) • Low molecular weight heparin (LMWH)² • LDUH or LMWH² combined with IPC or GCS
General Surgery Appendix A, Table 5.19	Any of the following: <ul style="list-style-type: none"> • Low-dose unfractionated heparin (LDUH) • Low molecular weight heparin (LMWH) • Factor Xa Inhibitor (Fondaparinux) • LDUH or LMWH or Factor Xa Inhibitor (fondaparinux) combined with IPC or GCS
General Surgery with a reason for not administering pharmacological prophylaxis Appendix A, Table 5.19	Any of the following: <ul style="list-style-type: none"> • Graduated Compression stockings (GCS) • Intermittent pneumatic compression devices (IPC)
Gynecologic Surgery Appendix A, Table 5.20	Any of the following: <ul style="list-style-type: none"> • Low-dose unfractionated heparin (LDUH) • Low molecular weight heparin (LMWH) • Factor Xa Inhibitor (fondaparinux) • Intermittent pneumatic compression devices (IPC) • LDUH or LMWH or Factor Xa Inhibitor (fondaparinux) combined with IPC or GCS
Urologic Surgery Appendix A, Table 5.21	Any of the following: <ul style="list-style-type: none"> • Low-dose unfractionated heparin (LDUH) • Low molecular weight heparin (LMWH) • Factor Xa Inhibitor (fondaparinux) • Intermittent pneumatic compression devices (IPC) • Graduated compression stockings (GCS) • LDUH or LMWH or Factor Xa Inhibitor (fondaparinux) combined with IPC or GCS
Elective Total Hip Replacement Appendix A, Table 5.22	Any of the following started within 24 hours of surgery: <ul style="list-style-type: none"> • Low molecular weight heparin (LMWH) • Factor Xa Inhibitor (Fondaparinux) • Warfarin